

Mathematics ECAT Pre Engineering Chapter 16 Solution of Trigonometric Equations Online Test

Sr	Questions	Answers Choice
1	Question Image	D. none of these
2	The solution of the equation $3 \tan^2 x = 1$ is _____	D. none of these
3	The solution set of the equation $4 \cos^2 x - 3 + 0$ is	D. none of these
4	The solution set of the equation $1 + \cos x = 0$ is _____	D. none of these
5	Question Image	D. none of these
6	Question Image	D. none of these
7	Question Image	D. none of these
8	Question Image	
9	Question Image	
10	Question Image	
11	Question Image	D. none of these
12	Question Image	A. I andII andI quadrants B. IIII andI andI andI quadrants C. IIII andI andI andI quadrants

13

Question Image

Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);>I and<span
style="color: rgb(34, 34, 34); font-
family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);>I<span style="color:
rgb(34, 34, 34); font-family:
"Times New Roman"; font-
size: 18px; background-color:
rgb(255, 255, 248);>V<span
style="color: rgb(34, 34, 34); font-
family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);>quadrants
D. none of these

A. <div><span style="color: rgb(34,
34, 34); font-family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);>I and<span
style="background-color: rgb(255,
255, 248); color: rgb(34, 34, 34);
font-family: "Times New
Roman"; font-size:
18px;">I<span
style="background-color: rgb(255,
255, 248); color: rgb(34, 34, 34);
font-family: "Times New
Roman"; font-size: 18px;">I
quadrants</div>
B. <span style="color: rgb(34, 34,
34); font-family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);><span style="color:
rgb(34, 34, 34); font-family:
"Times New Roman"; font-
size: 18px; background-color:
rgb(255, 255, 248);>I and
<span style="color: rgb(34, 34, 34);
font-family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);>I<span style="color:
rgb(34, 34, 34); font-family:
"Times New Roman"; font-
size: 18px; background-color:
rgb(255, 255, 248);>I quadrants
C. <span style="color: rgb(34, 34,
34); font-family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);><span style="color:
rgb(34, 34, 34); font-family:
"Times New Roman"; font-
size: 18px; background-color:
rgb(255, 255, 248);>I and
<span style="color: rgb(34, 34, 34);
font-family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);><span style="color:
rgb(34, 34, 34); font-family:
"Times New Roman"; font-
size: 18px; background-color:
rgb(255, 255, 248);>V<span
style="color: rgb(34, 34, 34); font-
family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);>quadrants
D. none of these

14

The solution set of $\sin x + \cos x = 0$ is

A. one element

B. two elements

C. three elements

D. Infinite elements

15

The solution set of trigonometric equation contains

16

General solution of $1 + \cos x = 0$ is

10 General solution of $1 + \cos x = 0$ is

17 Question Image D. all

18 Question Image

19 Question Image

20 Question Image

21 Question Image D. both a & c

22 Question Image

23 Question Image A. trigonometric equation
B. conditional equation
C. identity
D. None

24 For Cosine Rule of any triangle ABC, b^2 is equal to

A. $a^2 + b^2 - 2ab \cos C$
B. $a^2 + b^2 + 2ab \cos C$
C. $a^2 + b^2 + 2ab \cos B$
D. $a^2 + b^2 - 2ab \cos A$

25 In a triangle ABC, if angle A = 72° , angle B = 48° and c = 9 cm then C is

A. 69°
B. 66°
C. 60°
D. 63°

26 Considering Cosine Rule of any triangle ABC, possible measures of angle A includes

A. Angle A is obtuse
B. Angle A is acute
C. Angle A is right-angle
D. All of above

27 Sine rule for a triangle states that

A. $a/\sin A = b/\sin B = c/\sin C$
B. $\sin A/a = \sin B/b = \sin C/c$
C. $a/\sin A + b/\sin B + c/\sin C$
D. $2a/\sin A = 2b/\sin B = 2c/\sin C$

28 By expressing $\sin 125^\circ$ in terms of trigonometrical ratios, answer will be

A. $\sin 65^\circ = 0.9128$
B. $\sin 55^\circ = 0.8192$
C. $\sin 125^\circ = 0.8192$

- 0.95em;">sin 70° = 0.5384
- D. sin 72° = 0.1982
-
- 29 By expressing cos 113° in terms of trigonometrical ratios, answer will be
- A. cos 76° = -0.7093
B. cos 65° = -0.4258
C. cos 67° = -0.3907
D. cos 62° = -0.8520
-
- 30 Question Image
- 31 If Sin A = sin B, cos A = cos B, then the value of A in terms of B is
- 32 The general solution of $\tan 3x = 1$ is
- A. 30°
B. 45°
C. 60°
D. 75°
-
- 33 Question Image
- 34 If $4 \sin^2 \theta = 1$, then values of θ are
- A. No solution
B. One real solution
C. More than one real solution
D. None of these
-
- 35 Question Image
- 36 Question Image
- 37 Question Image
- 38 $\cot \theta = \sin 2\theta$ if $\theta =$
- 39 $\cot \theta = \sin 2\theta$ if $\theta =$
- 40 Question Image
- 41 Question Image
- A. 0
B. 5
C. 6
D. 10
-
- 42 The number of values of x in the interval $[0, 5\pi]$ satisfying the equation $3 \sin^2 x - 7 \sin x + 2 = 0$ is
- A. 0
B. 5
C. 6
D. 10
-
- 43 If $\sin 6\theta + \sin 4\theta + \sin 2\theta$, then $\theta =$
- A. 0
B. 1
C. 2
D. 3
-
- 44 The number of solution of the equation $\tan x + \sec x = 2 \cos x$ lying in the interval $[0, 2\pi]$ is
- A. A finite non-empty set
B. Null set
C. Both a and b
D. None of these
-
- 45 Question Image
- 46 The smallest positive root of the equation $\tan x - x = 0$ lies on
- 47 General solution of $\tan 5\theta = \cot 2\theta$ is
- 48 One root of the equation $\cos x - x + 1/2 = 0$ lies in the interval
- 49 The solution of the equation $\cos^2 \theta + \sin \theta + 1 = 0$ lies in the interval
- 50 If $\sin(\pi \cos \theta) = \cos(\pi \sin \theta)$, then which of the following is correct?
- A. 7
B. 5
C. 6
D. None of these
-
- 51 Question Image
- 52 Question Image
- A. From an empty set
B. 1
C. 2
D. >2
-
- 53 The general value of θ satisfying the equation $2 \sin^2 \theta - 3 \sin \theta - 2 = 0$ is

54	Question Image	
55	Question Image	A. 1 B. 2 C. 3 D. None of these
56	The number of points of intersection of two curves $y = 2 \sin x$ and $y = 5x^2 + 2x + 3$ is	A. 0 B. 1 C. 2 D. None of these
57	Question Image	D. none of these
58	The solution of the equation $3 \tan^2 x = 1$ is _____	D. none of these
59	The solution set of the equation $4 \cos^2 x - 3 + 0$ is	D. none of these
60	The solution set of the equation $1 + \cos x = 0$ is _____	D. none of these
61	Question Image	D. none of these
62	Question Image	D. none of these
63	Question Image	D. none of these
64	Question Image	
65	Question Image	
66	Question Image	
67	Question Image	D. none of these
68	Question Image	A. l andll andl quadrants B. ll andll andl quadrants C. ll andll andl quadrants D. ll andll andl quadrants

style="color: rgb(34, 34, 34); font-family: "Times New Roman"; font-size: 18px; background-color: rgb(255, 255, 248);">">Vquadrants

D. none of these

A. <div>I andI quadrants</div>
B. I andI quadrants

rgb(255, 255, 248);">I and
<span style="color: rgb(34, 34, 34);
font-family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);">I<span style="color:
rgb(34, 34, 34); font-family:
"Times New Roman"; font-size:
18px; background-color:
rgb(255, 255, 248);">I<span
style="color: rgb(34, 34, 34); font-
family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);">I quadrants
C. <span style="color: rgb(34, 34,
34); font-family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);">I<span style="color:
rgb(34, 34, 34); font-family:
"Times New Roman"; font-size:
18px; background-color:
rgb(255, 255, 248);">I and
<span style="color: rgb(34, 34, 34);
font-family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);">I<span style="color:
rgb(34, 34, 34); font-family:
"Times New Roman"; font-size:
18px; background-color:
rgb(255, 255, 248);">V<span
style="color: rgb(34, 34, 34); font-
family: "Times New
Roman"; font-size: 18px;
background-color: rgb(255, 255,
248);">quadrants
D. none of these

69

Question Image

70

The solution set of $\sin x + \cos x = 0$ is

71

The solution set of trigonometric equation contains

1

73

Question Image

A. one element

B. two elements

C. three elements

D. Infinite elements

Dall

		D. all
74	Question Image	
75	Question Image	
76	Question Image	
77	Question Image	D. both a & c
78	Question Image	
79	Question Image	A. trigonometric equation B. conditional equation C. identity D. None
80	For Cosine Rule of any triangle ABC, b^2 is equal to	<p>A. $a^2 - c^2 - 2ab \cos A$</p> <p>B. $a^2 + c^2 - 2ab \cos A$</p> <p>C. $a^2 + b^2 - 2ab \cos C$</p> <p>D. $a^2 + b^2 + 2ab \cos C$</p>
81	In a triangle ABC, if angle A = 72° , angle B = 48° and c = 9 cm then C is	<p>A. 69°</p> <p>B. 66°</p> <p>C. 60°</p> <p>D. 63°</p>
82	Considering Cosine Rule of any triangle ABC, possible measures of angle A includes	<p>A. Angle A is obtuse</p> <p>B. Angle A is acute</p> <p>C. Angle A is right-angle</p> <p>D. All of above</p>
83	Sine rule for a triangle states that	<p>A. $a/\sin A = b/\sin B = c/\sin C$</p> <p>B. $\sin A/a = \sin B/b = \sin C/c$</p> <p>C. $a/\sin A + b/\sin B + c/\sin C$</p> <p>D. $2a/\sin A = 2b/\sin B = 2c/\sin C$</p>
84	By expressing $\sin 125^\circ$ in terms of trigonometrical ratios, answer will be	<p>A. $\sin 65^\circ = 0.9128$</p> <p>B. $\sin 55^\circ = 0.8192$</p> <p>C. $\sin 70^\circ = 0.5384$</p> <p>D. $\sin 72^\circ = 0.982$</p>

85 By expressing $\cos 113^\circ$ in terms of trigonometrical ratios, answer will be

- A. $\cos 76^\circ = -0.7093$
- B. $\cos 65^\circ = -0.4258$
- C. $\cos 67^\circ = -0.3907$
- D. $\cos 62^\circ = -0.8520$

86 Question Image

87 If $\sin A = \sin B$, $\cos A = \cos B$, then the value of A in terms of B is

88 The general solution of $\tan 3x = 1$ is

- A. 30°
- B. 45°
- C. 60°
- D. 75°

90 If $4 \sin^2 \theta = 1$, then values of θ are

- A. No solution
- B. One real solution
- C. More than one real solution
- D. None of these

91 Question Image

92 Question Image

93 Question Image

94 $\cot \theta = \sin 2\theta$ if $\theta =$

95 $\cot \theta = \sin 2\theta$ if $\theta =$

96 Question Image

97 Question Image

98 The number of values of x in the interval $[0, 5\pi]$ satisfying the equation $3 \sin^2 x - 7 \sin x + 2 = 0$ is

- A. 0
- B. 5
- C. 6
- D. 10

99 If $\sin 6\theta + \sin 4\theta + \sin 2\theta = 0$, then $\theta =$

100 The number of solution of the equation $\tan x + \sec x = 2 \cos x$ lying in the interval $[0, 2\pi]$ is

- A. 0
- B. 1
- C. 2
- D. 3

101 Question Image

- A. A finite non-empty set
- B. Null set
- C. Both a and b
- D. None of these

102 The smallest positive root of the equation $\tan x - x = 0$ lies on

103 General solution of $\tan 5\theta = \cot 2\theta$ is

104 One root of the equation $\cos x - x + 1/2 = 0$ lies in the interval

105 The solution of the equation $\cos^2 \theta + \sin \theta + 1 = 0$ lies in the interval

106 If $\sin(\pi \cos \theta) = \cos(\pi \sin \theta)$, then which of the following is correct?

- A. 7
- B. 5
- C. 6
- D. None of these

108 Question Image

- A. From an empty set
- B. 1
- C. 2
- D. >2

109 The general value of θ satisfying the equation $2 \sin^2 \theta - 3 \sin \theta - 2 = 0$ is

110 Question Image

111 Question Image

- A. 1
- B. 2
- C. 3
- D. None of these

112 The number of points of intersection of two curves $y = 2 \sin x$ and $y = 5x^2 + 2x + 3$ is

- A. 0
- B. 1
- C. 2
- D. None of these