

## ECAT Mathematics Chapter 18 Basic Concepts & Definitions Online Test

Sr	Questions	Answers Choice
1	If $x = 1 - t^2$ and $y = 3t^2 - 2t^3$ then $dy/dx =$	A. $(1-t)$ B. $3(1+t)$ <b>C. <math>3(t-1)</math></b> D. $3/1-t$
2	Question Image	
3	Question Image	A. $-2x$ B. $x^{<sup>-3</sup>}$ D. $-2x^{<sup>3</sup>}$
4	If $y = x^m$ then $dy/dx$ equals:	A. $mx$ B. $x/m$ <b>C. <math>mx^{&lt;sup&gt;m-1&lt;/sup&gt;}</math></b> D. $x^{<sup>m-1</sup>}$
5	Question Image	<b>A. <math>2x</math></b> B. $x/2$ C. $2x^{<sup>3</sup>}$ D. $x^{<sup>3</sup>}/2$
6	If $f(x) = 1/x-2$ then $f^{-1}(0)$ equals:	<b>A. <math>-1/4</math></b> B. $-3/2$ C. $-1/2$ D. $1/5$
7	If $f(x) = x^{100}$ the value of $f^{-1}(1)$ is:	A. 100 <b>B. -100</b> C. 0 D. -101
8	Question Image	
9	$f(x) = ax^2 - 3x - 5$ , and $f^{-1}(2) = 9$ , a is equal to	<b>A. 2</b> B. 3 C. -2 D. 4
10	Question Image	
11	Question Image	A. zero at x <b>B. differentiable at x</b> C. continuous at x D. none of these
12	Question Image	
13	if $x \in D_f$ and $f^{-1}(x)$ exists, then $f$ is said to be	A. zero at x <b>B. Differentiable at x</b> C. Continuous at x D. None of these
14	If $3x + 4y + 7 = 0$ , then $dy / dx =$	A. $<div>-1/2</div>$ B. $-4/3$ C. $7/2$ <b>D. <math>-3/4</math></b>
15	Question Image	D. None of these
16	Question Image	
17	Question Image	<b>A. 100</b> B. -100 C. 0 D. -101
18	The derivative of $\sqrt{x}$ at $x = a$ is:	A. $1/2a$ B. $2 / \sqrt{a}$ C. $2\sqrt{x}$ <b>D. <math>1 / 2\sqrt{x}</math></b>
19	Question Image	
20	Question Image	

- 21  $d/dx(x^3 + 2x + 3) =$
- A.  $x^{<sup>2</sup>} + 2$   
 B.  $3x + 2$   
 C.  $3x^{<sup>2</sup>} + 5$   
 D.  $3x^2 + 2$

- 22 Question Image
- A.  $mx$   
 B.  $x/m$   
 C.  $mx^{<sup>m-1</sup>}$   
 D.  $xm^{<sup>m-1</sup>}$

- 23  $d/dx [\tan^2 x]$
- A.  $2\tan x \sec^2 x$   
 B.  $2\tan x \sec x$   
 C.  $2 \cot x \tan x$   
 D.  $2\sec^2 x \cos x$

- 24 If  $y = (7x + 9)^2$ , then  $dy/dx$  equals:
- A.  $98x + 126$   
 B.  $14x$   
 C.  $14x + 18$   
 D.  $14x + 81$

- 25 Question Image
- A.  $x^{<sup>2</sup>} + 2$   
 B.  $3x + 2$   
 C.  $3x^{<sup>2</sup>} + 5$   
 D.  $3x^{<sup>2</sup>} + 2$

- 27 If  $x^2 + y^2 = 1$ , then  $dy/dx$
- A.  $y/x$   
 B.  $-x/y$   
 C.  $1/x$   
 D. None of these

- 28 Question Image
- A. 8  
 B.  $1/8$   
 C.  $1/3$   
 D.  $2/3$

- 30 Question Image
- A. 0  
 B. 8  
 C. 5  
 D. 9

- 32  $d/dx (\operatorname{cosec} x)$
- A.  $-\sec x \tan x$   
 B.  $\sin x \cos x$   
 C.  $-\csc x \cot x$   
 D.  $2\sin x \cos x$

- 33 Question Image
- A. 1  
 B. 0  
 C.  $cx$   
 D. c

- 35 Question Image
- 36 Question Image

- 37 If  $y = 3x + 2\cos x$ , then  $dy/dx =$
- A.  $3-2 \sin x$   
 B.  $3-t \sin x$   
 C.  $3x^{<sup>2</sup>} - 2\sin x$   
 D.  $3(1-4 \sin x)$

- 38 If  $y = x^n$  then  $dy/dx$  equals:
- A.  $nx$   
 B.  $x^{<sup>n-1</sup>}$   
 C.  $nx^{<sup>n-1</sup>}$   
 D. n

- 39 Question Image
- 40  $d/dx (\cos x^2) =$
- A.  $-2x \cos x$   
 B.  $-2x \sin x^{<sup>2</sup>}$   
 C.  $-2x \tan x$   
 D.  $-2x \sec^2 x$

- 41 If  $f(x) = c$  then  $f^{-1}(x)$  equals:
- A. 1  
 B. 0  
 C.  $cx$   
 D. c

- 42 Question Image
- 43 Question Image B.  $x^{n-1}$
- 44 If  $f(x) = 2x^3 + 1$  then  $f'(0) =$  A. 0  
B. 1  
C. 6  
D. None of these
- 45 Question Image
- 46  $d/dx (\cos x \sin x) =$  A.  $\cos^2 x - \sin^2 x$   
B.  $2\cos^2 x + \sin^2 x$   
C.  $2\cos^2 x - \sin^2 x$   
D.  $1 - \sin^2 x$
- 47 Differentiation of  $\sin x$  w.r.t.  $\cot x$  is: A.  $-\sin^2 x \sec x$   
B.  $-\cos x \sin^2 x$   
C.  $-\cos^2 x \tan x$   
D.  $-\sin^2 x$
- 48  $d/dx (\cot x) =$  A.  $\sec x \tan x$   
B.  $-\csc^2 x$   
C.  $\sec^2 x$   
D.  $1/\cot^2 x$
- 49 If  $f(x) = x^{2/3}$  then  $f'(x)$  at  $x = 8$  equals: A. 8  
B.  $1/8$   
C.  $1/3$   
D.  $2/3$
- 50 Question Image
- 51 Let  $f$  be real valued function continuous in the interval  $(x, x_1) \subseteq D_f$  (the domain of  $f$ ), then  $f(x_1) - f(x)/x_1 - x$  represents: A. Instantaneous rate  
B. Average rate of change  
C. Differential coefficient  
D. None of these
- 52 If  $c$  is a constant, then  $d/dx (c) =$  A. 0  
B.  $c$   
C.  $cx$   
D. 1
- 53 Question Image
- 54 If  $f(x) = x^5 + x^3 + x$  the value of  $f'(1)$  is: A. 0  
B. 8  
C. 5  
D. 9
- 55 If  $y = 1/x^2$  then  $dy/dx$  equals: A.  $-2x$   
B.  $x^{-3}$   
C.  $-2/x^3$   
D.  $-2x^{-3}$
- 56 if  $y=x^2$  then  $dy/dx$  equals: A.  $2x$   
B.  $x/2$   
C.  $2x^3$   
D.  $x^3/2$
- 57 If  $x = at^2$  and  $y = 2at$  then  $dy/dx =$  A.  $2a/y$   
B.  $y/2a$   
C.  $-a/2y$   
D.  $-2y/a$
- 58 The derivative of  $1/x^m$  is: A.  $x^{m+1}/m$   
B.  $m(x)^{m-1}$   
C.  $(m-1)x^{-m}$   
D.  $m/x^{m+1}$
- 59 If  $f(x) = c$  then  $f'(x)$  equals: A. 1  
B. 0  
C.  $cx$   
D.  $c$
- 60 Question Image