

ECAT Mathematics Chapter 7 Partial Fractions Online Test

Sr	Questions	Answers Choice
1	Question Image	A. Improper rational fraction B. Proper rational fraction C. Polynomial D. Equation
2	Question Image	A. Polynomial B. Equation C. Improper rational fraction D. Proper rational fraction
3	$x^3 + 2x^2 - 3x + 5$ is _____	A. An equation B. A polynomial C. Proper rational fractions D. Improper rational fractions
4	$x^2 + x - 6 = 0$ is	A. An equation B. An identity C. A polynomial D. None of these
5	An open sentences formed by using the sign of equality '=' is called _____	A. An identity B. An equation C. A polynomial D. None of these
6	Question Image	
7	Question Image	
8	Question Image	
9	Question Image	
10	A fraction in which the degree of the numerator is less than the degree of the denominator is called	A. Polynomial B. Equation C. Proper fraction D. Improper fraction
11	Question Image	A. An expression B. Rational fraction C. Equation D. Identity
12	$(x + 3)(x + 4) = x^2 + 7x + 12$ is _____	A. Quadratic equation B. Linear equation C. Cubic equation D. Identity
13	Question Image	
14	Question Image	
15	Question Image	
16	Question Image	
17	Question Image	
18	A relation in which the equality is true only for some values of the known is called _____	A. An identity B. An equation C. A polynomial D. None of these
19	A relation in which the equality is true for all values of the unknown is called _____	A. An identity B. An equation C. A polynomial D. None of these
20	A fraction in which the degree of the numerator is greater than or equal to the degree of the denominator is called	A. A proper fraction B. An improper fraction C. An equation D. An identity

21	Question Image	
22	Question Image	
23	Question Image	
24	Question Image	
25	Question Image	
26	Question Image	
27	Question Image	
28	Question Image	
29	How many types of an equation	A. 1 B. 3 C. 2 D. None
30	An equation which holds good for all values of variables is called	A. Equation B. Conditional equation C. Constant D. None
31	An open sentence formed by using the sign of equality "=" is called	A. Equation B. In equation C. True sentence D. False sentence
32	$2x = 3$ is a conditional equation it is true for	A. 2 B. 3 C. $\frac{3}{2}$ D. $\frac{2}{3}$
33	Which is the proper rational function	
34	Question Image	A. $A = x, B = 1$ B. $A = 0, B = 2$ C. $A = -1, B = 1$ D. $A = x-1, B = x+1$
35	Question Image	
36	$(x + 2)^2 = x^2 + 4x + 4$ is	A. A linear equation B. A cubic equation C. A quadratic equation D. None
37	$x^2 + x - 6 = 0$ is a conditional equation and it is true for	A. 2, 3 B. 2, -3 C. -2, -3 D. -2, 3
38	The symbol _____ shall be used both for equation and identity	A.
39	Question Image	A. Improper rational fraction B. Rational fraction C. Proper rational fraction D. None of above
40	Question Image	A. Proper fraction B. Improper fraction C. Rational fraction D. None of these
41	Question Image	A. Rational fraction B. Proper fraction C. Improper rational fraction D. None of these
42	There are _____ types of rational fraction	A. Three B. Four C. Five D. Two
43	Question Image	
44	Which is a proper rational fraction	
45	Question Image	A. $A = x, B = 1$ B. $A = 0, B = 2$ C. $A = -1, B = 1$ D. $A = x-1, B = x + 1$

46	Question Image	
47	$(x + 2)^2 = x^2 + 4x + 4$ is	A. A linear equation B. A cubic equation C. A quadratic equation D. None
48	$x^2 + x - 5 = 0$ is	A. A polynomial B. An inequality C. An identity D. None
49	Question Image	
50	A fraction in which the degree of the numerator is less the degree of the denominator is called	A. Polynomial B. Proper fraction C. Rational fraction D. None
51	A relation in which the equality is true only for some values of the unknown is called	A. An identity B. An equation C. A polynomial D. None
52	Question Image	
53	When rational fraction is separated into partial fractions, the result is	A. an identity B. A fraction C. A partial sum D. Improper fraction
54	An improper rational fraction can be reduced by division to a	A. Proper fraction B. Polynomial C. mixed form
55	To express a single rational fraction as a sum of two or more single rational fractions which are called	A. improper fractions B. Partial fractions C. mixed form D. Polynomials
56	An equation which hold good for all values of the variables is called	A. Identity B. fraction C. mixed form D. Partial equation
57	Question Image	A. Improper rational fraction B. Proper rational fraction C. Polynomial D. Equation
58	Question Image	A. Polynomial B. Equation C. Improper rational fraction D. Proper rational fraction
59	$x^3 + 2x^2 - 3x + 5$ is _____	A. An equation B. A polynomial C. Proper rational fractions D. Improper rational fractions
60	$x^2 + x - 6 = 0$ is	A. An equation B. An identity C. A polynomial D. None of these
61	An open sentences formed by using the sign of equality '=' is called _____	A. An identity B. An equation C. A polynomial D. None of these
62	Question Image	
63	Question Image	
64	Question Image	
65	Question Image	
66	A fraction in which the degree of the numerator is less than the degree of the denominator is called	A. Polynomial B. Equation C. Proper fraction D. Improper fraction
67	Question Image	A. An expression B. Rational fraction C. Equation D. Identiv

68	$(x + 3)(x + 4) = x^2 + 7x + 12$ is _____	A. Quadratic equation B. Linear equation C. Cubic equation D. Identity
69	Question Image	
70	Question Image	
71	Question Image	
72	Question Image	
73	Question Image	
74	A relation in which the equality is true only for some values of the known is called _____	A. An identity B. An equation C. A polynomial D. None of these
75	A relation in which the equality is true for all values of the unknown is called _____	A. An identity B. An equation C. A polynomial D. None of these
76	A fraction in which the degree of the numerator is greater than or equal to the degree of the denominator is called	A. A proper fraction B. An improper fraction C. An equation D. An identity
77	Question Image	
78	Question Image	
79	Question Image	
80	Question Image	
81	Question Image	
82	Question Image	
83	Question Image	
84	Question Image	
85	How many types of an equation	A. 1 B. 3 C. 2 D. None
86	An equation which holds good for all values of variables is called	A. Equation B. Conditional equation C. Constant D. None
87	An open sentence formed by using the sign of equality "=" is called	A. Equation B. In equation C. True sentence D. False sentence
88	$2x = 3$ is a conditional equation it is true for	A. 2 B. 3 C. $\frac{3}{2}$ D. $\frac{2}{3}$
89	Which is the proper rational function	
90	Question Image	A. $A = x, B = 1$ B. $A = 0, B = 2$ C. $A = -1, B = 1$ D. $A = x-1, B = x+1$
91	Question Image	
92	$(x + 2)^2 = x^2 + 4x + 4$ is	A. A linear equation B. A cubic equation C. A quadratic equation D. None
93	$x^2 + x - 6 = 0$ is a conditional equation and it is true for	A. 2, 3 B. 2, -3 C. -2, -3 D. -2, 3

94	The symbol _____ shall be used both for equation and identity	A.
95	Question Image	A. Improper rational fraction B. Rational fraction C. Proper rational fraction D. None of above
96	Question Image	A. Proper fraction B. Improper fraction C. Rational fraction D. None of these
97	Question Image	A. Rational fraction B. Proper fraction C. Improper rational fraction D. None of these
98	There are _____ types of rational fraction	A. Three B. Four C. Five D. Two
99	Question Image	
100	Which is a proper rational fraction	
101	Question Image	A. $A = x, B = 1$ B. $A = 0, B = 2$ C. $A = -1, B = 1$ D. $A = x-1, B = x+1$
102	Question Image	
103	$(x+2)^2 = x^2 + 4x + 4$ is	A. A linear equation B. A cubic equation C. A quadratic equation D. None
104	$x^2 + x - 5 = 0$ is	A. A polynomial B. An inequality C. An identity D. None
105	Question Image	
106	A fraction in which the degree of the numerator is less the degree of the denominator is called	A. Polynomial B. Proper fraction C. Rational fraction D. None
107	A relation in which the equality is true only for some values of the unknown is called	A. An identity B. An equation C. A polynomial D. None
108	Question Image	
109	When rational fraction is separated into partial fractions, the result is	A. an identity B. A fraction C. A partial sum D. Improper fraction
110	An improper rational fraction can be reduced by division to a	A. Proper fraction B. Polynomial C. mixed form
111	To express a single rational fraction as a sum of two or more single rational fractions which are called	A. improper fractions B. Partial fractions C. mixed form D. Polynomials
112	An equation which hold good for all values of the variables is called	A. Identity B. fraction C. mixed form D. Partial equation