ECAT Physics Chapter 2 Vectors and Equilibrium

Sr	Questions	Answers Choice
1	Two forces each of the magnitude F act perpendicular to each other. The angle made by the resultant force with the horizontal will be:	A. 30<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° B. 45<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° C. 60<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° D. 90<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">°
2	Torque is also called:	A. Momentum B. Linear inertia C. Moment of a force D. Mass
3	The resultant of two velocities 3 m/sec and 400 cm/sec making an angle 90° with each other is:	A. 20 m/sec B. 5 m/sec C. 3 m.sec D. None of these
4	The vector in space has:	A. One component B. Two components C. Three components D. None of these
5	All trigonometric functions (sine, cosine, tangent etc) are positive in:	A. 1st quadrant B. 2nd quadrant C. 3rd quadrant D. 4th quadrant
6	If a vector lies in second quadrant, than $B_x$and $B_y$are:	A. -,+ B. +,- C. +,+ D. -,-
7	When a vector is multiplied by a negative number, its direction:	A. Remains the same B. Changes C. Changes by 180<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° D. None of these

8	Tick the correct answer:	A. Torque is a vector quantity B. Torque is the turning effect of a force C. Torque is called moment of a force D. All of above
9	When the magnitude of two component vectors are equal to that of their resultant, then the angle between the components is:	A. 60<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° B. 90<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° C. 120<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° D. 150<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">°
10	Two forces each of 10 N act on a body, if the force are inclined at 30° and 60° respectively with x-axis, then x-component of their resultant is:	A. 20 N B. 13.66 N C. 10 N D. 8.66 N
11	A person starts his journey from a point 0, travels 4 Km SW, then 4 Km NW, and finally 4 Km north-east. At what distance is he now from point 0?	A. 0 Km B. 4 Km C. 8 Km D. 12 Km
12	The magnitude of resultant of three vectors is 3. Its x-component is one, y-component is two, then its z-component is:	A. 0 B. 1 C. 2 D. 3
13	A force of 5 n is acting Y-axis. Its component along X-axis is:	A. 7 N B. 5 N C. Zero D. 10 N
14	The rectangular components of a vector are equal in magnitude when the vector makes and angle _____with their x-component:	A. 0<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° B. 30<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° C. 45<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">°

		D. 60<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">°
15	The magnitude of the resultant of two forces may be increased by:	A. Increasing the angle between them B. Decreasing the angle between them C. Drawing a triangle to represent them D. None of these
16	The direction of vector in space is specified by:	A. One angle B. Two angles C. Three angles D. None of these
17	By convention, torques producing clockwise rotation are taken as:	A. Positive B. Nagative C. Zero D. None of these
18	If the vector 5 N lies along with x-axis, then its component along y-axis will be:	A. Zero B. 5 N C. 7 N D. 10 N
19	Two vectors to be combined have magnitudes of 60 N and 35 N. Pick the possible answer:	A. 100 N B. 70 N C. 20 N D. Zero
20	Two vectors having different magnitudes:	A. Have their directions opposite B. May have their resultant zero C. Cannot have their resultant zero D. None of these