

Physics ECAT Pre Engineering Chapter 20 Atomic Spectra Physics Online Test

Sr	Questions	Answers Choice
1	The transitions of electrons in the hydrogen atom result in the emission of spectral lines in the:	A. Ultra red region B. Visible region C. Ultraviolet region D. Any of these E. None of these
2	The natural arrangement of colours in the spectrum of white light spectrum is	A. VIBGYOR B. ROYBGIV C. ROYBIGV D. BIGROYV E. None of these
3	Lyman series in the spectrum of hydrogen exists in the :	A. Infra-red region B. Visible region C. Ultraviolet region D. Both (A) and (B) E. None of these
4	Static electricity is produced by the transfer of:	A. Electrons B. Protons C. One fluid D. Two fluid E. None of these
5	The first series which was identified in the spectrum of hydrogen is called:	A. Lyman series B. Balmer series C. Paschen series D. Brackett series E. Pfund series
6	Spectrum represents the number of component colours present in certain light in terms of:	A. Wavelength B. Frequency C. Energy D. Both (A) and (B) E. All of these
7	The value of the metastable state for Neon is	A. 20.66eV B. 20.61eV C. 19.23eV D. 18.70eV
8	Tick the series which lie/s in the infra-red region.	A. Pfund series B. Brackett series C. Paschen series D. All of these E. None of these
9	Graph of Black body radiation is example of	A. Band spectra B. Continuous spectra C. Line spectra D. All
10	An electron of the hydrogen atom in the second orbit is called its:	A. Ground state B. Excited state C. Ionized state D. Any of these E. None of these
11	The minimum wavelength of X-rays produced of 1KV potential difference is applied across the anode and cathode of the tube is	A. 1.24×10^{-10} m B. 7.92×10^{-20} m C. 2.78×10^{-14} m D. 3.88×10^{-11} m
12	X-ray are also known as	A. Roentgen rays B. Maxwell rays C. Plank range D. Einstein rays
13	Selenium is:	A. An insulator B. A conductor C. Both A and B D. Excellent conductor E. None of these
		A. L-shell

14	The first shell near the nucleus is	B. X-shell C. N-shell D. M-shell
15	The results of spectra obtained by Balmer were expressed in 1896 by	A. Bohr B. Rydberg C. Planck D. Rutherford E. Coulomb
16	The lasing or active medium in He-Ne laser discharge tube is:	A. Nitrogen B. Helium C. Hydrogen D. Neon E. None of these
17	The shell closer to the nucleus is called:	A. N shell B. L shell C. K shell D. M shell E. O shell
18	Laser is a beam of:	A. Visible light B. Infra red light C. Ultra violet light D. Violet light only E. yellow light only
19	The process of formation of spectrum is called:	A. Interference B. Spectroscopy C. Dispersion D. Reflection E. Both (A) and (D)
20	Photons must have energy equal to	A. ev B. En C. hf D. None of these
21	CT scanning is the abbreviated name of	A. Computed Technology B. Computed Technique C. Computed Technology D. Computerized Technique
22	Atoms of hydrogen gas can be excited by passing electric current through it when the gas is filled into the discharge tube at a pressure which is	A. Less than atmospheric pressure B. Much less than atmospheric pressure C. Greater than atmospheric pressure D. Much greater than atmospheric pressure E. Both C and D
23	If the distance between two charges is doubled, the force between them will become:	A. Double B. Half C. One third D. One fourth
24	Balmer series was identified in:	A. 1685 B. 1785 C. 1885 D. 1985 E. 1585
25	The photocopying process is called:	A. Geography B. Sonography C. Xerography D. Zerography E. None of these
26	Balmer series lies in that region of electromagnetic wave spectrum which is called:	A. Visible region B. Invisible region C. Infra-red region D. ultraviolet region E. None of these
27	The results of spectra obtained by Balmer were expressed in 1896 by:	A. Bohr B. Rydberg C. Planck D. Rutherford E. Coulomb
28	The range of wavelengths of colours in the visible colours is	A. 140 nm to 456 nm B. 10 nm to 56 nm C. 410 nm to 656 nm D. 910 nm to 956 nm E. None of these

29	The formula of Brackett series can be obtained by putting in the general formula, the value of n equal to:	<p>A. four</p> <p>B. two</p> <p>C. three</p> <p>D. four</p> <p>E. five</p>
30	An compared to solid matter, a crack or an air bubble allows:	<p>A. Great amount of X-rays to pass</p> <p>B. Smallest amount of X-rays to pass</p> <p>C. Very small amount of X-rays to pass</p> <p>D. Any of these</p> <p>E. None of these</p>
31	The electric field lines start from:	<p>A. Positive charge</p> <p>B. Negative charge</p> <p>C. Either A and B</p> <p>D. Neutron</p> <p>E. An atom</p>
32	A metastable state:	<p>A. Is an excited state</p> <p>B. Is that in which excited electron is stable</p> <p>C. Is that in which excited electron is usually unstable</p> <p>D. Means a time interval of 10^{-8} second</p> <p>E. Both (A) and (C)</p>
33	In flesh, light element like carbon, hydrogen and oxygen predominate. Three elements allows _____ amount of incident X-ray to pass through them	<p>A. Small</p> <p>B. Greater</p> <p>C. Equal</p> <p>D. Sometimes</p>
34	Coulomb multiplied by volt by volt gives the unit called:	<p>A. farad</p> <p>B. Ohm</p> <p>C. Second</p> <p>D. joule</p> <p>E. Watt</p>
35	The holes created in the L and M shells are occupied by transitions of:	<p>A. Electrons from lower states</p> <p>B. Electrons from higher state</p> <p>C. Positrons from higher states</p> <p>D. Electrons from K shell</p> <p>E. Both (A) and (B)</p>
36	In helium Neon Laser Neon = 15% and Helium = 85% used. The lasing gas this unit is	<p>A. Helium</p> <p>B. Neon</p> <p>C. Both</p> <p>D. None of these</p>
37	Tick the series which lies in the visible region:	<p>A. Lyman series</p> <p>B. Balmer series</p> <p>C. Paschen series</p> <p>D. Brackett series</p> <p>E. P fund series</p>
38	The inkjet printer eject a thin stream of:	<p>A. Water</p> <p>B. Oil</p> <p>C. Ink</p> <p>D. Any above</p> <p>E. None of these</p>
39	By CAT scans, we can detect the density difference of the order of:	<p>A. 1%</p> <p>B. 20%</p> <p>C. 30%</p> <p>D. 50%</p> <p>E. 70%</p>
40	In case of braking radiations, when the rate of deceleration is very large, the emitted radiation corresponds to:	<p>A. Short wavelength</p> <p>B. Large wavelength</p> <p>C. Very large wavelength</p> <p>D. Low frequency</p> <p>E. Both (B) and (C)</p>
41	Energy required by an electron revolving in certain orbit to jump to an excited state is called:	<p>A. Ionization energy</p> <p>B. Ionization potential</p> <p>C. Excitation energy</p> <p>D. Excitation potential</p> <p>E. None of these</p>
42	The life time of metastable state is equal to	<p>A. Life time of excited state</p> <p>B. Greater than by excited state</p> <p>C. Zero</p> <p>D. Less than by excited state</p>
43	Gaussian surface is always:	<p>A. Rectangular</p> <p>B. Spherical</p> <p>C. Cylinder</p> <p>D. Box shape</p>

		E. Any of these
44	The He-Ne laser discharge tube is filled with:	A. 85% He B. 15% He C. 50% He D. 60% He E. 85% Ne
45	The spectrum emitted from hydrogen filled discharge tube is:	A. Line spectrum B. Discrete spectrum C. And spectrum D. Absorption spectrum E. Both (A) and (B)
46	Braking radiation causes:	A. Continuous spectrum B. Line Spectrum C. Band spectrum D. Discrete spectrum E. All of these
47	Ultraviolet region lies in _____ series	A. Lyman B. Balmer C. P fund D. B racket
48	Consider a photon of continuous X-ray and a photon of characteristics X-ray of same wavelength. Which of the following is/are different for the two photons	A. Frequency B. Penetrating power C. Energy D. Method of creation
49	X-rays produced in a tube operating at 10^5 V. The speed of X-rays produced is	A. 3×10^8 m/s B. 3.1×10^8 m/s C. 2.8×10^8 m/s D. 1.88×10^8 m/s
50	X-rays can penetrate in a solid matte through a distance of several:	A. Kilo metres B. Metres C. Centimeters D. A few angstroms E. One micrometer
51	We can excite an atom by	A. Bombardment of particles B. Radiating photons C. Providing potential difference D. All answer are true