



neight="19"  
src="file:///C:/Users/Softsol/AppData/Local/Temp/msohtmlclip1/01/clip\_image002.png"  
v:shapes="\_x0000\_i1025" style="font-family: Calibri, sans-serif; font-size: 11pt;"><!--  
[endif--></span><!--[endif--><o:p></o:p></p>

13	The geometric mean of the a observations 2,4,8, is:	A. 2 B. 8 C. 4 D. no geometric mean
14	The formula of range is:	A. $X_{\max} - X_{\min}$ B. $X_{\max} + X_{\min}$ C. groups/wight&nbsp;nbsp; D. none of these
15	Venn diagram was first used by.....	A. John Venn B. Netwon C. Arthur Cayler D. John Napier
16	A and A <sup>c</sup> are.....Set.	A. Universal B. Overlapping C. Disjoint D. Super
17	A collection of well-defined distinct object is called:	A. Subset B. Power set C. Set D. None of these
18	The different number of ways to describe a set are:	A. 1 B. 2 C. 3 D. 4
19	A set containing no element is called.	A. subset B. Empty set C. Singleton set D. Super set
20	If $x \in U$ and $x \notin A$ , then $\{x\}$ is equal to .....	A. $U^c$ B. $A^c$ C. $\emptyset^c$ D. $A - U$