

Chemistry Fsc Part 1 Chapter 8 Online Test

Sr	Questions	Answers Choice
1	A buffer solution can be prepared by mixing	A. Weak acid and its salt with weak base B. Weak base and its salt with strong acid C. Strong acid and its salts with weak base D. Strong base and its salt with weak acid
2	The pK_a value of CH_3COOH is 4.74 when we mix CH_3COOH and CH_3COONa in the ratio of 10:1, then the pH of the buffer is	A. 4.74 B. 5.74 C. 3.74 D. 7.00
3	The sum of pH and pOH is	A. 0 B. 7 C. 14 D. 10
4	A solution has pH zero. Its H^+ ions concentration will	A. zero B. More than unity C. Less than unity D. Unity only
5	Which acid has less value of pK_a .	A. CH_3COOH B. H_2S C. H_2CO_3 D. HCl
6	Which statement is correct about solubility product constant.	A. It is applicable at highly soluble substances. B. Value of K_{sp} is independent of temperature C. It is used for homogeneous aquarium system D. It can be used to predict that precipitation will take place or not by combining two ions
7	For which system does the equilibrium constant K_c has units of $(\text{concentration})^{-1}$	
8	K_w for water at 0 °C is 0.1×10^{-34} and at 100 °C 7.5×10^{-14} , How many times dissociation of water increase from 0 °C to 100 °C	A. 7.5 times B. 50 times C. 75 times D. 100 times
9	Question Image	A. $\text{dm}^3 \times 10^{-6} \times \text{mole}^{-2}$ B. $\text{mole}^2 \times \text{dm}^{-6}$ C. Mole dm^{-3} D. Having no units
10	Catalyst used in preparation of NH_3 from N_2 and H_2 is.	A. Ni B. Fe C. Pt D. V_2O_5
11	pH of buffer is calculated by.	A. Sorenson equation B. Mosley equation C. Henderson equation D. De broglie equation
12	_____ was derived by C.M Guldberg and P Waage in 1864	A. Law of conservation of Mass B. Law of mass action C. Law of conservation of energy D. Distribution law
13	Buffer action can be explained by	A. Common ion effect B. Law of mass action C. Le Chateller's principle D. All above
14	The relationship between K_p and K_c is given by	

15	A solution will be unsaturated if	A. Ionic product = K_{sp} B. Ionic product $< K_{sp}$ C. Ionic Product $> K_{sp}$ D. both 'a' and 'b' are correct
16	The number of moles of acid or base required by one dm^3 of buffer to alter its pH by one unit is called	A. Buffer efficiency B. Buffer capacity C. Buffer action D. None
17	When HCl is added to H_2S aqueous solution, its ionization	A. Decrease B. Increase C. Remains constant D. First increases than decreases
18	The effect of temperature on equilibrium was studied by	A. Lewis B. Van der waal C. Arrhenius D. Vant hoff
19	An excess of aqueous silver nitrate is added to aqueous barium chloride and precipitate is removed by filtration. What are the main ions in the filtrate	
20	pH of rain water.	A. 7 B. Slightly basic C. slightly acidic D. Highly basic
21	When small amount of acid or base is added to buffer, its pH.	A. Remain same B. Always increases C. Always decreases D. slightly increases or decreases
22	The solubility product of AgCl is $2.0 \times 10^{-10} \text{ mol}^2 \text{ dm}^{-6}$ The maximum concentration of Ag^+ ions in the solution is	A. $2.0 \times 10^{-10} \text{ mol dm}^{-3}$ B. $1.41 \times 10^{-5} \text{ mol dm}^{-3}$ C. $1.0 \times 10^{-10} \text{ mol dm}^{-3}$ D. $4.0 \times 10^{-20} \text{ mol dm}^{-3}$
23	The pH of human blood is	A. 7.0 B. 7.4 C. 4.0 D. 6.5
24	Some impurities of MgCl_2 are present in NaCl which separation technique can be used to separate the impurities.	A. Filtration B. Crystallization C. Common ion effect D. Chromatography
25	Which one of the following aqueous solutions has the highest pH	A. 0.1 M NaOH B. 0.1 M HCl C. 0.2 M H_2SO_4 D. 0.1 M HNO_3
26	The law of mass action was given by	A. D.C. down and P wage B. Gay Lussic and C.M C. C.M Goldberg and P. Waage D. Hendeson and Le Chateller's
27	Optimum pressure in Haber's process for synthesis of Ammonia is	A. 100 -150 atm B. 200- 300 atm C. 350 - 450 atm D. 500 - 600 atm
28	pK_a of CH_3COOH is 4.74. The pK_b value of CH_3COO^- ions will be	A. 7 B. 14 C. 9.26 D. zero
29	One dm^3 of a buffer solution containing 0.01 M NH_4Cl and 0.1 M NH_4OH having pK_a of 3 has pH.	A. 4 B. 6 C. 9 D. 10
30	K_a and K_b of a conjugate acid and are related with K_w as	A. $K_a \times K_b = K_w$ B. $K_a \times K_b = K_w$ C. $K_a \times K_b = K_w$ D. $K_a \times K_b = K_w$

A. Decreasing the temperature

31	For the equilibrium system $\text{N}_2 + \text{O}_2 + \text{Heat} = 2\text{NO}$ the equilibrium constant decreases by	B. Adding a catalyst C. Adding N_2 D. Adding NO
32	A solution have H^+ ions concentration 1×10^{-7} , its pH will	A. Acidic B. Basic C. Neutral D. Zero
33	The pH of 10^{-3} mole dm^{-3} of an aqueous solution of H_2SO_4 is.	A. 3.0 B. 2.7 C. 2.0 D. 1.5
34	the substance which increase the rate of reaction but remains unchanged at the end of the reaction is called.	A. Indicator B. Promoter C. Catalyst D. Activated complex
35	Almost forward reaction is complete when value of K_c is	A. very high B. Very small C. Neither large nor very small D. No correlation
36	The unit of K_c for the reaction $\text{N}_2 + \text{O}_2 = 2\text{NO}$ will be	A. mol dm^{-3} B. $\text{mol}^{-1} \text{ dm}^3$ C. $\text{mol}^{-2} \text{ dm}^6$ D. No units
37	The pH of $10^{-3} \text{ mol dm}^{-3}$ of an aqueous solution of H_2SO_4 is	A. 3.0 B. 2.7 C. 2.0 D. 1.5
38	When solid KI dissolved in water, its heat of solution is positive. What would happen to dissolution when temperature is increased.	A. Increases B. Decreases C. Remain same D. Firs increases than decreases
39	Sum of pK_a and pK_b is equal to.	A. 1 B. 7 C. 0 D. 14
40	Question Image	A. The value of K_{p} falls with a rise in temperature B. The value of K_{p} falls with increasing pressure C. Adding V_2O_5 catalyst increase the equilibrium yield of sulphur trioxide D. The value of K_{p} is equal to K_{c}
41	If the volumes of reactants and products are same in a gaseous phase reaction, then the equilibrium state is not affected by	A. Change of temperature B. Change of pressure C. Change of concentration D. Catalyst
42	When concentration of one product is removed at equilibrium stage, in which direction it moves to reestablish equilibrium.	A. Forward B. Reverse C. Neither forward nor reverse D. Equally move in both direction
43	The value of pH and pOH of pure water at 25°C is	A. 14 B. 7 C. 1×10^{-14} D. 1×10^{14}
44	Le-Chatelier Braun principle is sometimes known as	A. Law of mass action B. Law of mobile equilibrium C. Law of active mass D. All of these above
45	Which one of the following has highest pH	A. Distilled water B. $1 \text{ M NH}_4\text{OH}$ C. 1 M NaOH D. Water saturated with chlorine gas
46	At equilibrium stage of chemical reaction	A. The concentration of reaction is equal to concentration of products B. The rate constant of forward reaction is equal to rate constant of backward reaction C. The rate of forward reaction is equal rate of backward reaction D. The energy of activation of forward step is equal to energy of activation of

47 A chemical reaction $A \rightleftharpoons B$ is said to be in equilibrium when

- A. Complete conversion of A to B has taken place
- B. Conversion of A to B is 50% complete
- C. Rate of transformation of A to B is equal to B to A
- D. 50% Reactant have been changed to B