

Business Mathematics Icom Part 1 Chapter 4 Online Test

| 0 | | A 01 : |
|----|---|---|
| Sr | Questions | Answers Choice |
| 1 | The sign of every equation is: | A. ≠ B. = C. <u>></u> D. <u><</u> |
| 2 | Solution set of $4x - 7y = 12$ and $3x + y = 9$ is | A. (0,3) B. (1,3) C. (6,3) D. (3,0) |
| 3 | Both sides of an equation are joined by | A. > B. < C. = D. ≠ |
| 4 | 1 : 3 is same as: | A. 3 to 1 B. 3 : 8 C. 1 to 3 D. None of the above |
| 5 | Two consecutive odd integers are: | A. x and (x + 2) B. (x + 1) and (x + 3) C. 2x, (2x + 2) D. (2x + 1) and (2x + 3) |
| 6 | If $3^{2x} + a = 10 \ 3^{x}$ in transformed from is $y^{2} + 9 = 10y$, then the transformation is: | A. 3 ^{2x} = y B. 3 ^x = y C. 1/3 ^x = y D. None of these |
| 7 | Aa ^x + Ba ^{-x} = C is a standard form of | A. Exponential equationB. Linear equationC. Quadratic equationD. Reciprocal equation |
| 8 | The solution set of equation $x^2 + 2x + 1 = 0$ is | A. {1} B. {-1} C. {1, -1} D. None of these |
| 9 | The power of variable in a quadratic equation is | A. 3 B. 1 C. 4 D. 2 |
| 10 | A set of simultaneous equation is called set of inconsistent equation if: | A. Value of one of the unknown obtained B. Value of one of the unknown obtained C. Values of all the unknown obtained D. None of these |
| 11 | Simultaneous equations can be solved in ways. | A. 2 B. 3 C. 4 D. 5 |
| 12 | Formula to calculate compounded amount is: | A. P(1 + i) ⁿ B. P(1 + i) ⁻ⁿ C. R (1 + i) D. P (1 - i) ⁿ |
| 13 | Solution set of equations $4x + 5y = 40$ and $3x + 2y = 23$ is: | A. { (4,5) } B. { (5, 4) } C. { (-5, 4) } D. { -4, -5) } |
| 14 | 90.5% in common fraction: | A. 0.9 B. 10/9 C. 9/10 |

| | | 门ノ | |
|--|--|----|--|
| | | | |
| | | | |

| | | D. 101/200 |
|----|---|---|
| 15 | Equation of the form $ax^4 + bx^3 + bx + a$ is: | A. Polynomial B. Reciprocal C. Irrational D. None of these |
| 16 | A linear equation always has: | A. Three roots B. Two roots C. One root D. No root |
| 17 | Factorization is one of the method use to solve: | A. $ax + b = 0$ B. $ax2 + bx + C = 0$ C. $ax3 + bx + c = 0$ D. None of these |
| 18 | 5x - 2 = 10 is a | A. Open sentence B. Right sentence C. False sentence D. Equation |
| 19 | In quadratic equation the variable has degree: | A. 1 B. 2 C. More than 2 D. Less than 2 |
| 20 | The solution set for a quadratic equation x^2 -8x + 15 is | A. (3, 5) B. (-3, -5) C. (3, -5) D. (-5, 3) |
| 21 | A linear equation consist of roots | A. One B. Two C. Zero D. Three |
| 22 | ${\sf B}^2$ - 4ac in a quadratic formula is called | A. Nature of rootB. DiscriminantC. Solution setD. Extraneous root |
| 23 | System of simultaneous equations is solved by: | A. FactorizationB. Subtraction of additionC. SubstitutionD. Both b and c |
| 24 | The roots of quadratic equation will be imaginary if b^2 - 4ac is | A. 0 Bve C. +ve D. Greater than zero |
| | | |