

## Mathematics English Medium 5th Class Online Test

Sr	Questions	Answers Choice
1	In 856211, the place value of ... is ten thousands.	A. 6 B. 5 C. 8 D. 7
2	$209856 - 205660 =$	A. 9149 B. 9641 C. 6419 D. 4196
3	If a price of motorcycle is 79459, then price of 2 motorcycles will be.	A. 148918 B. 150918 C. 152918 D. 153918
4	A quadrilateral having only one pair of parallel sides is called.	A. trapezium B. Square C. riombus D. traingle
5	Cabe is a .....solid figure.	A. Two dimensoal B. Three dimentional C. Four dimentional D. Five dimentional
6	1 cm -----	A. 100 mm B. 10 mm C. 1/10 mm D. 1000 mm
7	To find the sum of given values whose average is knowns, the following formula is used.	A. sum of item = average of item + number of item B. sum of item= average of item-number C. sum of items = average of item x number of items D. None of these
8	0.5 will be witeen is percentage	A. 5% B. 0.5% C. 50% D. 500%
9	The greatest 6 -digit number is	A. 999999 B. 983007 C. 100000 D. 9,999999
10	The product of common prime factorization is called.	A. LCM B. Factorization C. HCF D. Factor
11	Percentage is a special hind of fraction with as denominator	A. 10 B. 100 C. 1000 D. 1
12	6 min =.....sec	A. 360 B. 1/60 C. 42 D. 180
13	The average price of soe books is Rs. 80. If total price of books is Rs.400, then total number of books will be.	A. 5 B. 6 C. 8 D. 10
14	The average of 4,6,8 in	A. 4 B. 6 C. 8 D. 10
		A. 1400 mm B. 1500 mm C. 1600 mm D. 1700 mm

15 the length of pencil is 14 cm . IN milimeteres it is.  
B. 140 mm  
C. 1.4 mm  
D. 14 mm

---

16 A triangle in which one angle is obtuse is called.  
A. Acute angled traingle  
B. Obtuse angled traingle  
C. Right angled traingle  
D. Equilaceral triangle

---

17 When the value of one item is known, the value of many item of sameof kind can be found by.  
A. Addition  
B. Subraction  
C. Multiplication  
D. Division

---

18 To convert milometeres into meteress, by 1000  
A. Add  
B. Subtrat  
C. Divide  
D. Multiply

---

19 A..... is a 2-dimensioal shape which can be folded in a specific pattern to get a 3-dimensioal solid.  
A. Net  
B. rhombus  
C. Triangle  
D. parrlelogram

---

20 When a numebr is divided by another number, the left over quantity is called.  
A. Quotient  
B. Remainder  
C. Addition  
D. Substraction