

## Business Statistics Icom Part 2 English Medium Online Test

Sr	Questions	Answers Choice
1	Questionnaire method is used in the collection of.	A. Constant B. Variable C. Parameter D. Statistic
2	Collection of data by village patwari an example of.	A. Secondary data B. Primary data C. Grouped data D. Non of above
3	Geometric mean of the relatives is.	A. Non -reversible B. Reversible C. Both a and b D. None of above
4	A card is drawn fram an ordinary pack of 52 cards. The probability that it is red, and either an ace or a heart is.	A. 2/52 B. 1/13 C. 1/52 D. 2/13
5	Data classified by attributes is called.	A. Quantitative B. Qualitative C. Numerical D. None of above
6	An event that contains more than one sample point is called.	A. Compound event B. Independent event C. Simple event D. Multiple event
7	In fixed base method the base period should be.	A. Normal year B. Abnormal year C. Fluctuating year D. Both b and c
8	Sample mean is	A. Variable B. Statistic C. Parameter D. Constant
9	The probability of drawing black cards from a pack of 52 cards is.	A. 13/52 B. 12/52 C. 26/52 D. 4/52
10	The six faces of the die are called equality likely if the die is.	A. Six -faced B. Round C. Fair D. Steeper
11	Coded method of calculation is nly used in	A. A.M B. Median C. Combined mean D. None of above
12	A measure computed on the basis of population data is called.	A. Parameter B. Statistics C. Statistic D. None of these
13	All elements in a particualr case represent.	A. Universe B. Sample C. Population D. Both a and c
14	The term bar means. a.	A. Thick wide line B. Thin wide line C. Thick narrow line
15	Lower class boundary of 30-35 will be.	A. 35 B. 36 C. 32.5 D. None of above

16	Graph of time series is also called.	A. Band graph B. Line graph <b>C. Range graph</b> D. None of above
17	Fisher index number is the G.M. of the.	A. Marshall Edgeworth index number <b>B. Laspeyres's and Paasche's index number</b> C. Laspeyres's index number D. Paasche's index number
18	The probability of an event cannot be.	A. More than one B. Less than one <b>C. Negative</b> D. Zero
19	As a general rule, statisticians tend to use which of the following number of classes when arranging the data.	<b>A. Between 5 and 20</b> B. Fewer than 5 C. Between 1 and 5 D. More than 20
20	In a symmetrical distribution, mean, median and mode are always.	A. Different B. Zero <b>C. Identical</b> D. Negative