Physics ECAT Pre Engineering Chapter 2 Vectors and Equilibrium

Sr	Questions	Answers Choice
1	All trigonometric functions (sine, cosine, tangent etc) are positive in:	A. 1st quadrant B. 2nd quadrant C. 3rd quadrant D. 4th quadrant
2	The magnitude of resultant of three vectors is 3. Its x-component is one, y-component is two, then its z-component is:	A. 0 B. 1 C. 2 D. 3
3	If x-component of a vector is -3 N and y-component is 3 N, then angle of resultant vector will x-axis is:	A. 45<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° B. 315<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° C. 135<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° D. 225<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">°
4	A person starts his journey from a point 0, travels 4 Km SW, then 4 Km NW, and finally 4 Km north-east. At what distance is he now from point 0?	A. 0 Km B. 4 Km C. 8 Km D. 12 Km
5	The vector is space has:	A. One Component B. Two Compnents C. Three Components D. Non of these
6	Tick the correct answer:	A. Torque is a vector quantity B. Torque is the turning effect of a force C. Torque is called moment of a force D. All of above
		A. 30<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° B. 45<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial,

7	Two forces each of the magnitude F act perpendicular to each other. The angle made by the resultant force with the horizontal will be:	sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° C. 60<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° D. 90<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">°
8	Two forces of 10 N and 8 N are applied simultaneously to a body. the maximum value of their resultant is:	A. 2 N B. - 2 N C. 18 N D. 36 N
9	The sum of two or more vectors is equal to a single vector which is called:	A. Component vector B. Resultant vector C. Product vector D. None of these
10	When a vector is multiplied by a negative number, its direction:	A. Remains the same B. Changes C. Changes by 180<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° D. None of these
11	Torque is also called:	A. Momentum B. Linear inertia C. Moment of a force D. Mass
12	Unit vector is used to specify:	A. Magnitude of a vector B. Dimensions of a vector C. Direction of a vector D. Position of a vector
13	An vector of 10 N makes an angle of 45° with x-axis. Angle between its rectangular components with be:	A. 45<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° B. 90<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° C. 135<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° D. Zero
		A. Zero B. 5 N

14	If the vector 5 N lies along with x-axis, then its component along y-axis will be:	B. 5 N C. 7 N D. 10 N
15	Two vectors having different magnitudes:	A. Have their directions opposite B. May have their resultant zero C. Cannot have their resultant zero D. None of these
16	Choose the set of physical quantities, which have both numerical and directional properties:	A. Velocity, mass B. Speed, acceleration C. acceleration weight D. Distance, force
17	The magnitude of the resultant of two forces may be increased by:	A. Increasing the angle between them B. Decreasing the angle between them C. Drawing a triangle to represent them D. None of these
18	A vector of magnitude 5 N is added to a vector of magnitude 8 N while the orientations are changeable. Range of their possible sum will be very from:	A. Zero to 3 N B. 1 N to 13 N C. 13 N to 3 N D. None of these
19	Parallel vectors of same magnitudes:	A. Are equal B. Are unequal C. When added give the some equal to zero D. Give the answer equal to zero
20	The vector in space has:	A. One component B. Two components C. Three components D. None of these