

Statistics Ics Part 1 Chapter 8 Online Test

| Sr | Questions | Answers Choice |
|----|--|---|
| 1 | Random number can be generated manually by | A. Drawing cards from numbered cards B. Roiting or spinning numbered wheels C. Use of random numbers table D. All of these |
| 2 | The probability of success changes from trial to trial, is the property of: | A. Binomial experiment B. Hypergeometric experiment C. Both A and B D. None of these |
| 3 | The probability of even/odd number when a fair die rolled is: | A. 1/6 B. 2/6 C. 1/36 D. 3/6 |
| 4 | The number of possible outcomes in Bernoulli trial is | A. Three B. Four C. Two D. One |
| 5 | If in binomial distribution, $\mu = 6$, $p = 3/5$, the number of trial are: | A. 18 B. 30 C. 10 D. None of these |
| 6 | Which of the following case is true for hypergeometric distribution. | A. Probability remains constant for all trials B. Probability changes C. successive trials are dependent D. Both (B) and (C) but not (A) |
| 7 | N-K is classified as | A. Success B. Failures C. Parameters D. Range |
| 8 | The expected value of a discrete random variable is. | A. Always an integer B. Always one of the values that the random variable can assume C. An interval of values D. None of these |
| 9 | The probability of success ----- from one trial to another when trials are dependent | A. Remains constant B. Is changed C. One D. Zero |
| 10 | If X and Y are independent random variables, the $E(XY)$ is equal to. | A. $E(XY)$ B. $E(X)E(Y)$ C. $XE(Y)$ D. $YE(X)$ |
| 11 | A fair die is rolled three times. The probability of getting three "aces" is | A. 1/3 B. 1/6 |
| 12 | Binomial distribution is used when n is | A. Large B. Small C. Negative D. Zero |
| 13 | The parameters of the binomial distributions are | A. x and n B. x and p C. p and q D. n and p |
| 14 | The mean of binomial distribution is always: | A. Equal to variance B. Less than variance C. Greater than variance D. None of these |
| 15 | The parameters of hypergeometric distribution are: | A. n, k & p B. n, k & q C. n, p & q D. n, k & q |

| | | |
|----|--|---|
| 16 | If $E(X) = E$ then find arithmetic means will be. | A. 1 B. 4 C. 0 D. 8 |
| 17 | The repeated trials of binomial experiments are | A. Dependent B. Independent C. Fixed D. Variable |
| 18 | Binomial distribution is positively skewed | A. $P \leq 1/2$ B. $P \geq 1/2$ C. $P = 1/2$ D. $P \leq q$ & $q \leq P$ |
| 19 | If the random variable X denotes the number of heads when three distinct coins are tossed, then X assumes the value. | A. 0,1,2,3 B. 1,3,3,1 C. 1,2,3 D. None of these |
| 20 | In binomial distribution, the random variable has a range: | A. 0,1,2,...,n B. 0,1,2,...,+ ∞ C. - ∞ to + ∞ D. ∞ to +0 |